

BOILING POINT ELEVATION PROBLEMS

(nonelectrolytes)

Table of K_b values for solvents

Solvent	K_b ($^{\circ}\text{C/molal}$)	Solvent	K_b ($^{\circ}\text{C/molal}$)
Water	0.52	Benzene	2.67
Ethanol	1.20	Carbon tetrachloride	5.02
Chloroform	3.85	Acetic Acid	2.93

1. What is the boiling point elevation when 11.4 g of ammonia (NH_3) is dissolved in 200 g of H_2O ?
 2. How many grams of benzoic acid ($\text{C}_7\text{H}_6\text{O}_2$) must be dissolved in 78.1 g of ethanol to raise the boiling point by 4.00°C ?
 3. If 67.7 g of urea ($\text{CH}_4\text{N}_2\text{O}$) is dissolved in 833 g of chloroform, what is the elevation in boiling point?
 4. How many grams of camphor ($\text{C}_{10}\text{H}_{16}\text{O}$) are needed to raise the boiling point of 43.5 g of benzene by 2.10°C ?
 5. If 1800 g of ethylene glycol is added to 1900 g of water, what is the elevation in boiling point?
 6. If the boiling point of 69.6 g of carbon tetrachloride must be raised by 10.2°C , how many grams of pyridine ($\text{C}_5\text{H}_5\text{N}$) must be dissolved in the carbon tetrachloride?
 7. What is the boiling point elevation when 31.5 g of menthol ($\text{C}_{10}\text{H}_{20}\text{O}$) is dissolved in 258 g of acetic acid?
 8. How much will the boiling point of 25.0 g of acetic acid be raised if 2.69 g of picolinic acid ($\text{C}_6\text{H}_5\text{N}_2$) is dissolved in the acetic acid?
 9. Styrene glycol ($\text{C}_8\text{H}_{10}\text{O}_2$) is a plasticizer. How many grams of styrene glycol must be dissolved in 98.7 g of benzene to raise the boiling point by 8.57°C ?
 10. What is the boiling point elevation when 43.5 g of the dye magenta-I ($\text{C}_{20}\text{H}_{20}\text{ClN}_3$) is dissolved in 1650 g of ethanol?
-