

PERSPECTIVE

## Novel Features of G Protein-Coupled Receptor Kinase 4

Kim A. Neve

*Department of Behavioral Neuroscience, Oregon Health & Science University; and Veterans Affairs Medical Center, Portland, Oregon*

Received December 7, 2005; accepted December 9, 2005

### ABSTRACT

The defining characteristic of G protein-coupled receptor homologous desensitization is that the receptor must be occupied by an agonist or in an activated conformation that mimics an agonist-induced state. In most instances, the mechanistic basis for this characteristic is the high selectivity of G protein-coupled receptor kinases for the activated receptor. In this issue, Rankin et al. (p. 759) demonstrate that under some conditions, at least, the G protein-coupled receptor kinase GRK4 does not display a preference for the agonist-occupied D<sub>1</sub> dopamine receptor. Coexpression of GRK4 and the D<sub>1</sub> receptor in a heterologous

system induces phosphorylation of the receptor in the absence of agonist, causing constitutive desensitization and internalization of the receptor. Lacking the normal rapid feedback mechanisms associated with homologous desensitization, a system incorporating constitutively active GRK4 will be prone to dysregulation, perhaps explaining the generally low expression of GRK4. Indeed, considerable evidence suggests that just such dysregulation resulting from mutationally activated GRK4 contributes to the heritable component of human essential hypertension (*Physiol Genomics* **19**:223–246, 2004).

G protein-coupled receptors (GPCRs) typically respond to agonist stimulation with a time-dependent, rapidly reversible diminution, or desensitization, of the signaling response. Desensitization can be homologous (a receptor desensitized as a result of its own activation) or heterologous (one receptor desensitized as a consequence of the activation of and signaling by a different receptor). The canonical model of homologous desensitization of GPCRs (Van Koppen and Jakobs, 2004) is that the agonist-activated receptor binds and activates a GPCR-selective kinase (GRK) (Benovic et al., 1986), which phosphorylates the receptor on multiple serine/threonine residues. Activation and phosphorylation of the GPCR increases its affinity for arrestin (Benovic et al., 1987). Binding of arrestin to the intracellular loops of the GPCR both sterically hinders the interaction of receptor and G protein and recruits the GPCR into clathrin-coated pits for dynamin-dependent internalization into clathrin-coated vesicles (Goodman et al., 1996). The internalized GPCR may be dephosphorylated and rapidly recycled to the plasma mem-

brane (Pippig et al., 1995) or retained in the cell and ultimately degraded, with the choice determined by factors such as the phosphorylation state of other residues (Mason et al., 2002) and the stability of the interaction between the GPCR and arrestin (Oakley et al., 1999; Pan et al., 2003).

This model of homologous desensitization has been validated for countless GPCRs. Still, at almost every step of this process, there are examples of exceptions to and deviations from the model. In some cases, the deviations represent expanded roles for some of the players, such as GRKs (Fig. 1); in other cases, the deviations represent alternative pathways in addition to those in the canonical pathway. In still other cases, it seems that a receptor uses an alternative mechanism instead of the canonical pathway. The GRK can bind to the receptor and influence signaling without phosphorylating the receptor (Perroy et al., 2003; Dhami et al., 2005), GRKs phosphorylate other proteins constitutively or upon activation by a GPCR (Pitcher et al., 1998b; Hall et al., 1999; Pronin et al., 2000), arrestin can bind to unphosphorylated receptor (Chen et al., 2004; Kim et al., 2004; Jala et al., 2005), GPCR-activated arrestin can mediate GPCR signaling in addition to internalization (Luttrell and Lefkowitz, 2002; Gurevich and Gurevich, 2003), desensitization and internal-

Article, publication date, and citation information can be found at <http://molpharm.aspetjournals.org>.

doi:10.1124/mol.105.021535.

Please see the related article on page 759.

**ABBREVIATIONS:** GPCR, G protein-coupled receptor; GRK, G protein-coupled receptor kinase;  $\beta$ ARK,  $\beta$ -adrenergic receptor kinase; HEK, human embryonic kidney.

ization can occur without phosphorylation of the GPCR (Malcz et al., 1998) or without arrestin binding (Pals-Rylaarsdam et al., 1997; Bennett et al., 2001; Bhatnagar et al., 2001; Van Koppen and Jakobs, 2004), GPCRs can internalize via clathrin- and dynamin-independent pathways (Pals-Rylaarsdam et al., 1997; Vickery and von Zastrow, 1999), and dephosphorylation/resensitization can occur in the plasma membrane without receptor internalization (Gardner et al., 2001).

One component of the model that has been constant, and that defines homologous desensitization, has been the strong preference of GRKs for phosphorylation of the agonist-activated receptor over the inactive receptor. The instances when a GPCR is robustly phosphorylated by a GRK without being occupied by an agonist are examples that prove the rule, because the receptor invariably has a high level of inherent or mutationally induced constitutive activity (Ren et al., 1993; Pei et al., 1994; Geras-Raaka et al., 1998; Miller et al., 2003; Marion et al., 2004); the GPCR must be in an active conformation for significant GRK-catalyzed phosphorylation to occur. There is also some evidence that a GRK bound to an activated GPCR might phosphorylate adjacent, inactive GPCRs (Palczewski, 1997), but that does not invalidate the fundamental requirement for an activated receptor to bind and activate the GRK.


In this issue of *Molecular Pharmacology*, Rankin et al. (2006) demonstrate that heterologously expressed dopamine D<sub>1</sub> receptor is constitutively phosphorylated by heterolo-

gously expressed GRK4. The D<sub>1</sub> receptor does not have an unusually high level of constitutive activity, yet when coexpressed in HEK293 cells with GRK4, it is phosphorylated to such an extent that agonist treatment causes little additional phosphorylation; in this system, GRK4 apparently does not distinguish between active and inactive D<sub>1</sub> receptor. GRK4-catalyzed constitutive phosphorylation is associated with reduced dopamine-stimulated cyclic AMP accumulation (desensitization) and receptor internalization. The constitutively phosphorylated residues seem to be near the C terminus of the receptor, because truncation at residue Thr404 or combined mutation of Thr428 and Ser431 prevents or substantially decreases constitutive phosphorylation, desensitization, and internalization of the receptor. Thus, the inactive D<sub>1</sub> receptor can be phosphorylated by GRK4, resulting in diminished responsiveness to subsequent stimulation by dopamine.

The seven GRKs are grouped into three subfamilies: the retinal subfamily (GRK1/7), the  $\beta$ -adrenergic receptor kinase ( $\beta$ ARK) subfamily (GRK2/3), and the GRK4 subfamily (GRK4/5/6). Characteristics of the GRK4 subfamily include predominant localization at the membrane as a result of palmitoylation on C-terminal cysteine residues (for GRK4/6) or interaction between a positively charged domain near the C terminus and negatively charged membrane phospholipids (GRK5), activation by phosphatidylinositol bisphosphate binding to an N-terminal domain, and enhanced sensitivity to inhibition by calcium-sensor proteins such as calmodulin (Pronin et al., 1997; Pitcher et al., 1998a; Kohout and Lefkowitz, 2003; Willets et al., 2003). GRK4 is unusual within its subfamily (but similar to GRK1) in that its relatively low sequence homology across species suggests that it is subject to lower evolutionary pressure for sequence conservation and evolving more rapidly than the other members of its subfamily and the  $\beta$ ARK subfamily (Premont et al., 1999). It is interesting that GRK1/7 and GRK4 also differ from the other GRKs in tissue distribution. GRK2/3 and GRK5/6 are ubiquitously expressed, whereas GRK1/7 are expressed almost exclusively in the retina, and GRK4 is abundantly expressed only in the testes and expressed at much lower levels in other tissues including the kidney and the brain (Ambrose et al., 1992; Premont et al., 1996; Virilon et al., 1998; Salles et al., 2000; Willets et al., 2003).

Human GRK4 exists as four splice variants: GRK4 $\alpha$ , GRK4 $\beta$ , GRK4 $\gamma$ , and GRK4 $\delta$  (Premont et al., 1996). GRK4 $\alpha$  is the full-length version, most homologous with the other GRKs. GRK4 $\beta$  is missing the sequence encoded by exon 2, resulting in a 32-residue deletion that encompasses the phosphatidylinositol bisphosphate binding domain near the N terminus. GRK4 $\gamma$  is missing the sequence encoded by exon 15, resulting in a 46-residue deletion near the C terminus, and GRK4 $\delta$ , the shortest variant, is missing both alternatively spliced exons. Rankin et al. (2006) determined that the D<sub>1</sub> receptor was constitutively phosphorylated only by coexpression with GRK4 $\alpha$  and not with GRK2, GRK3, or any of the shorter splice variants of GRK4.

This work raises many interesting questions pertaining to the specificity of the response. First, is this a unique characteristic of the D<sub>1</sub> receptor, or will other GPCRs be found to be constitutively phosphorylated by GRK4? GRK4 is capable of phosphorylating and/or desensitizing activated forms of rhodopsin (Virilon et al., 1998), the follicle-stimulating hormone



**Fig. 1.** Expanded role for G protein-coupled receptor kinases in desensitization and signaling. 1, a GPCR is shown precoupled to a heterotrimeric G protein before activation by agonist. 2, in the canonical model of homologous desensitization, the agonist-activated GPCR binds to and is phosphorylated by GRK, which promotes the binding of arrestin, thus interfering with coupling to the G protein. 3, GRK binds to the receptor and noncatalytically hinders GPCR coupling to the G protein. Receptor-activated GRK also phosphorylates other proteins, such as tubulin. 4, Rankin et al. (2006) demonstrate that GRK4 constitutively phosphorylates a nonagonist-occupied GPCR, leading to desensitization and internalization, presumably via binding of arrestin. 5, GRK constitutively phosphorylates other receptor-interacting proteins, such as the Na<sup>+</sup>/H<sup>+</sup> exchanger regulatory factor (NHERF). 6, GRK2/3, in particular, may also decrease signaling by binding to G $\alpha$  and accelerating its GTPase activity (Willets et al., 2003), stimulating the hydrolysis of GTP to GDP and inorganic phosphate.

receptor (Lazari et al., 1999), the m2 muscarinic receptor (Tsuga et al., 1998), the luteinizing hormone/chorionic gonadotropin receptor (Premont et al., 1996), and the  $\beta_2$ -adrenoceptor (Premont et al., 1996). GRK4 can also regulate the calcium-sensing receptor (Pi et al., 2005) and is the endogenous GRK that mediates homologous desensitization of two other class C GPCRs in cerebellar neurons, the mGluR1 (Sallese et al., 2000) and GABA<sub>B</sub> (Perroy et al., 2003) receptors, in addition to being an endogenous regulator of the D<sub>1</sub> receptor in renal proximal tubule cells (Felder et al., 2002; Watanabe et al., 2002). There is one interesting report that GRK4 causes constitutive phosphorylation of the  $\beta_2$ -adrenoceptor in HEK293 cells, with no additional phosphorylation induced by agonist treatment, and also causes enhanced agonist-independent internalization of the receptor (Ménard et al., 1996). It seems likely that GRK4 will be found to catalyze constitutive phosphorylation of additional GPCRs.

A second question is whether constitutive GPCR phosphorylation is restricted to the GRK4 subtype. Rankin et al. (2006) determined that the D<sub>1</sub> receptor is not constitutively phosphorylated by GRK2/3, the two members of the  $\beta$ ARK subfamily. Although other members of the GRK4 subfamily were not tested by Rankin et al. (2006), previous work has shown robust agonist-stimulated phosphorylation of the D<sub>1</sub> receptor by GRK5 in HEK293 cells (Tiberi et al., 1996), suggesting a preference of that kinase for the activated state of the receptor. For the  $\beta_2$ -adrenoceptor, all three members of the GRK4 subfamily caused significantly more basal phosphorylation than GRK1–3, but only in the presence of GRK4 was there no additional agonist-induced phosphorylation (Ménard et al., 1996). A mechanistic basis for the greater propensity of members of the GRK4 subfamily to phosphorylate inactive GPCRs could be their constitutive localization at the membrane, in contrast to GRK2/3, whose translocation to the membrane is aided by free G $\beta\gamma$  produced by GPCR-activated heterotrimeric G proteins. If GRK4 is more likely than other members of that subfamily to exhibit no preference for activated GPCR over inactive receptor, an interesting line of investigation will be to identify the unique features of GRK4 that are responsible for this characteristic.

A system in which a GRK constitutively desensitizes a GPCR seems susceptible to dysregulation, in contrast to the homeostasis conferred by homologous desensitization in which only the activated receptor is desensitized, because an overabundance of the GRK or a mutation that enhances its activity, as in the kidney (see below), could cause perpetual desensitization. Is this the reason for the restricted distribution and generally low abundance of GRK4? Is there a unique characteristic of GPCR function in testes that makes it advantageous to have a high level of GRK4 and, hypothetically, constitutive desensitization?

Finally, what is the physiological relevance of the constitutive phosphorylation of the D<sub>1</sub> receptor by GRK4? Significant expression of GRK4 in D<sub>1</sub> receptor-dense brain regions has not been described. In renal proximal tubules, on the other hand, where the D<sub>1</sub> receptor regulates natriuresis, genetic hypertension in rats and human essential hypertension are associated with nonresponsiveness to dopamine because of constitutive desensitization of the D<sub>1</sub> receptor (Zeng et al., 2004). GRK4, in particular the GRK4 $\gamma$  splice variant, catalyzes D<sub>1</sub> receptor hyperphosphorylation in renal proximal tubule cells from subjects with essential hypertension

(Felder et al., 2002). Essential hypertension is linked to a locus that includes GRK4 (Casari et al., 1995) and is associated with nonsynonymous single nucleotide polymorphisms in the coding region of GRK4 (Speirs et al., 2004). When heterologously expressed with the D<sub>1</sub> receptor in Chinese hamster ovary cells, allelic variants of GRK4 $\gamma$  (R65L, A142V, A486V) cause enhanced desensitization and agonist-independent phosphorylation of the receptor, and transgenic mice expressing the A142V variant of GRK4 $\gamma$ , but not wild-type GRK4 $\gamma$ , are hypertensive and lack D<sub>1</sub> agonist-induced diuresis and natriuresis (Felder et al., 2002). The parallels between the work of Rankin et al. (2006) and the role of GRK4 in the kidney are not exact, because GRK4 $\gamma$  seems to be the variant that regulates the D<sub>1</sub> receptor in renal proximal tubule cells, whereas in HEK293 cells, GRK4 $\gamma$  and its allelic variants, as well as GRK4 $\beta$  and GRK4 $\delta$ , do not enhance basal phosphorylation of the D<sub>1</sub> receptor or have any effect that can be distinguished from endogenous GRKs (Rankin et al., 2006). It is possible that the lack of effect of GRK4 $\gamma$  in HEK293 cells can be attributed to differences in the cellular environment, and an interesting line of investigation will be to evaluate the effect of GRK4 $\alpha$ , and the effect of the single nucleotide polymorphisms in the context of GRK4 $\alpha$ , on D<sub>1</sub> receptor function in renal proximal tubule cells. Despite the discrepancies, the similarity between the observation that GRK4 does not distinguish between inactive and active D<sub>1</sub> receptor in HEK293 cells and the accumulating evidence that GRK4 hyperactivity is the cause of insensitivity to dopamine in essential hypertension suggests that further investigation of the mechanisms of this unusual characteristic of GRK4 will also help to elucidate fundamental mechanisms of the disorder.

## References

- Ambrose C, James M, Barnes G, Lin C, Bates G, Altherr M, Duyao M, Groot N, Church D, Wasmuth JJ, et al. (1992) A novel G protein-coupled receptor kinase gene cloned from 4p16.3. *Hum Mol Genet* **1**:697–703.
- Bennett TA, Foutz TD, Gurevich VV, Sklar LA, and Prossnitz ER (2001) Partial phosphorylation of the N-formyl peptide receptor inhibits G protein association independent of arrestin binding. *J Biol Chem* **276**:49195–49203.
- Benovic JL, Kühn H, Weyand I, Codina J, Caron MG, and Lefkowitz RJ (1987) Functional desensitization of the isolated  $\beta$ -adrenergic receptor by the  $\beta$ -adrenergic receptor kinase: potential role of an analog of the retinal protein arrestin (48-kDa protein). *Proc Natl Acad Sci USA* **84**:8879–8882.
- Benovic JL, Strasser RH, Caron MG, and Lefkowitz RJ (1986) Beta-adrenergic kinase: identification of a novel protein kinase that phosphorylates the agonist-occupied form of the receptor. *Proc Natl Acad Sci USA* **83**:2797–2801.
- Bhatnagar A, Willins DL, Gray JA, Woods J, Benovic JL, and Roth BL (2001) The dynamin-dependent, arrestin-independent internalization of 5-hydroxytryptamine 2A (5-HT<sub>2A</sub>) serotonin receptors reveals differential sorting of arrestins and 5-HT<sub>2A</sub> receptors during endocytosis. *J Biol Chem* **276**:8269–8277.
- Casari G, Barlassina C, Cusi D, Zagato L, Muirhead R, Righetti M, Nembri P, Amar K, Gatti M, Macciardi F, et al. (1995) Association of the alpha-adducin locus with essential hypertension. *Hypertension* **25**:320–326.
- Chen CH, Paing MM, and Trejo J (2004) Termination of protease-activated receptor-1 signaling by  $\beta$ -arrestins is independent of receptor phosphorylation. *J Biol Chem* **279**:10020–10031.
- Dhami GK, Babwah AV, Sterne-Marr R, and Ferguson SS (2005) Phosphorylation-independent regulation of metabotropic glutamate receptor 1 signaling requires G protein-coupled receptor kinase 2 binding to the second intracellular loop. *J Biol Chem* **280**:24420–24427.
- Felder RA, Sanada H, Xu J, Yu PY, Wang Z, Watanabe H, Asico LD, Wang W, Zheng S, Yamaguchi I, et al. (2002) G protein-coupled receptor kinase 4 gene variants in human essential hypertension. *Proc Natl Acad Sci USA* **99**:3872–3877.
- Gardner B, Liu ZF, Jiang D, and Sibley DR (2001) The role of phosphorylation/dephosphorylation in agonist-induced desensitization of D<sub>1</sub> dopamine receptor function: evidence for a novel pathway for receptor dephosphorylation. *Mol Pharmacol* **59**:310–321.
- Geras-Raaka E, Arvanitakis L, Bais C, Cesarman E, Mesri EA, and Gershengorn MC (1998) Inhibition of constitutive signaling of Kaposi's sarcoma-associated herpesvirus G protein-coupled receptor by protein kinases in mammalian cells in culture. *J Exp Med* **187**:801–806.
- Goodman OB Jr, Krupnick JG, Santini F, Gurevich VV, Penn RB, Gagnon AW, Keen JH, and Benovic JL (1996)  $\beta$ -Arrestin acts as a clathrin adaptor in endocytosis of the  $\beta_2$ -adrenergic receptor. *Nature (Lond)* **383**:447–450.

- Gurevich VV and Gurevich EV (2003) The new face of active receptor bound arrestin attracts new partners. *Structure* **11**:1037–1042.
- Hall RA, Spurney RF, Premont RT, Rahman N, Blitzer JT, Pitcher JA, and Lefkowitz RJ (1999) G protein-coupled receptor kinase 6A phosphorylates the Na<sup>+</sup>/H<sup>+</sup> exchanger regulatory factor via a PDZ domain-mediated interaction. *J Biol Chem* **274**:24328–24334.
- Jala VR, Shao WH, and Haribabu B (2005) Phosphorylation-independent  $\beta$ -arrestin translocation and internalization of leukotriene B<sub>4</sub> receptors. *J Biol Chem* **280**:4880–4887.
- Kim OJ, Gardner BR, Williams DB, Marinac PS, Cabrera DM, Peters JD, Mak CC, Kim KM, and Sibley DR (2004) The role of phosphorylation in D<sub>1</sub> dopamine receptor desensitization: evidence for a novel mechanism of arrestin association. *J Biol Chem* **279**:7999–8010.
- Kohout TA and Lefkowitz RJ (2003) Regulation of G protein-coupled receptor kinases and arrestins during receptor desensitization. *Mol Pharmacol* **63**:9–18.
- Lazari MFM, Liu X, Nakamura K, Benovic JL, and Ascoli M (1999) Role of G protein-coupled receptor kinases on the agonist-induced phosphorylation and internalization of the follitropin receptor. *Mol Endocrinol* **13**:866–878.
- Luttrell LM and Lefkowitz RJ (2002) The role of  $\beta$ -arrestins in the termination and transduction of G-protein-coupled receptor signals. *J Cell Sci* **115**:455–465.
- Malecz N, Bambino T, Bencsik M, and Nissenson RA (1998) Identification of phosphorylation sites in the G protein-coupled receptor for parathyroid hormone. Receptor phosphorylation is not required for agonist-induced internalization. *Mol Endocrinol* **12**:1846–1856.
- Marion S, Weiner DM, and Caron MG (2004) RNA editing induces variation in desensitization and trafficking of 5-hydroxytryptamine 2c receptor isoforms. *J Biol Chem* **279**:2945–2954.
- Mason JN, Kozell LB, and Neve KA (2002) Regulation of dopamine D<sub>1</sub> receptor trafficking by protein kinase A-dependent phosphorylation. *Mol Pharmacol* **61**:806–816.
- Ménard L, Ferguson SSG, Barak LS, Bertrand L, Premont RT, Colapietro AM, Lefkowitz RJ, and Caron MG (1996) Members of the G protein-coupled receptor kinase family that phosphorylate the  $\beta_2$ -adrenergic receptor facilitate sequestration. *Biochemistry* **35**:4155–4160.
- Miller WE, Houtz DA, Nelson CD, Kolattukudy PE, and Lefkowitz RJ (2003) G-protein-coupled receptor (GPCR) kinase phosphorylation and  $\beta$ -arrestin recruitment regulate the constitutive signaling activity of the human cytomegalovirus US28 GPCR. *J Biol Chem* **278**:21663–21671.
- Oakley RH, Laporte SA, Holt JA, Barak LS, and Caron MG (1999) Association of  $\beta$ -arrestin with G protein-coupled receptors during clathrin-mediated endocytosis dictates the profile of receptor resensitization. *J Biol Chem* **274**:32248–32257.
- Palczewski K (1997) GTP-binding-protein-coupled receptor kinases—two mechanistic models. *Eur J Biochem* **248**:261–269.
- Pals-Rylaarsdam R, Gurevich VV, Lee KB, Ptasiński JA, Benovic JL, and Hosey MM (1997) Internalization of the m2 muscarinic acetylcholine receptor: arrestin-independent and -dependent pathways. *J Biol Chem* **272**:23682–23689.
- Pan L, Gurevich EV, and Gurevich VV (2003) The nature of the arrestin-receptor complex determines the ultimate fate of the internalized receptor. *J Biol Chem* **278**:11623–11632.
- Pei G, Samama P, Lohse M, Wang M, Codina J, and Lefkowitz RJ (1994) A constitutively active mutant  $\beta_2$ -adrenergic receptor is constitutively desensitized and phosphorylated. *Proc Natl Acad Sci USA* **91**:2699–2702.
- Perroy J, Adam L, Qanbar R, Chenier S, and Bouvier M (2003) Phosphorylation-independent desensitization of GABA<sub>B</sub> receptor by GRK4. *EMBO (Eur Mol Biol Organ)* **J** **22**:3816–3824.
- Pi M, Oakley RH, Gesty-Palmer D, Cruickshank RD, Spurney RF, Luttrell LM, and Quarles LD (2005)  $\beta$ -Arrestin- and G protein receptor kinase-mediated calcium-sensing receptor desensitization. *Mol Endocrinol* **19**:1078–1087.
- Pippig S, Andexinger S, and Lohse MJ (1995) Sequestration and recycling of  $\beta_2$ -adrenergic receptors permit receptor resensitization. *Mol Pharmacol* **47**:666–676.
- Pitcher JA, Freedman NJ, and Lefkowitz RJ (1998a) G protein-coupled receptor kinases. *Annu Rev Biochem* **67**:653–692.
- Pitcher JA, Hall RA, Daaka Y, Zhang J, Ferguson SSG, Hester S, Miller S, Caron MG, Lefkowitz RJ, and Barak LS (1998b) The G protein-coupled receptor kinase 2 is a microtubule-associated protein kinase that phosphorylates tubulin. *J Biol Chem* **273**:12316–12324.
- Premont RT, Macrae AD, Aparicio SAJR, Kendall HE, Welch JE, and Lefkowitz RJ (1999) The GRK4 subfamily of G protein-coupled receptor kinases: alternative splicing, gene organization and sequence conservation. *J Biol Chem* **274**:29381–29389.
- Premont RT, Macrae AD, Stoffel RH, Chung N, Pitcher JA, Ambrose C, Ingles J, MacDonald ME, and Lefkowitz RJ (1996) Characterization of the G protein-coupled receptor kinase GRK4. Identification of four splice variants. *J Biol Chem* **271**:6403–6410.
- Pronin AN, Morris AJ, Surguchov A, and Benovic JL (2000) Synucleins are a novel class of substrates for G protein-coupled receptor kinases. *J Biol Chem* **275**:26515–26522.
- Pronin AN, Satpaev DK, Slepak VZ, and Benovic JL (1997) Regulation of G protein-coupled receptor kinases by calmodulin and localization of the calmodulin binding domain. *J Biol Chem* **272**:18273–18280.
- Rankin ML, Marinac PS, Cabrera DM, Wang Z, Jose PA, and Sibley DR (2006) The D<sub>1</sub> dopamine receptor is constitutively phosphorylated by G protein-coupled receptor kinase 4: evidence for a novel mechanism of regulation. *Mol Pharmacol* **69**:759–769.
- Ren Q, Kurose H, Lefkowitz RJ, and Cotecchia S (1993) Constitutively active mutants of the  $\alpha_2$ -adrenergic receptor. *J Biol Chem* **268**:16483–16487.
- Sallese M, Salvatore L, D'Urbano E, Sala G, Storto M, Launey T, Nicoletti F, Knöpfel T, and De Blasi A (2000) The G-protein-coupled receptor kinase GRK4 mediates homologous desensitization of metabotropic glutamate receptor 1. *FASEB J* **14**:2569–2580.
- Speirs HJ, Katyk K, Kumar NN, Benjafeld AV, Wang WY, and Morris BJ (2004) Association of G-protein-coupled receptor kinase 4 haplotypes, but not *HSD3B1* or *PTPIB* polymorphisms, with essential hypertension. *J Hypertens* **22**:931–936.
- Tiberi M, Nash SR, Bertrand L, Lefkowitz RJ, and Caron MG (1996) Differential regulation of dopamine D1A receptor responsiveness by various G protein-coupled receptor kinases. *J Biol Chem* **271**:3771–3778.
- Tsuga H, Okuno E, Kameyama K, and Haga T (1998) Sequestration of human muscarinic acetylcholine receptor hm1-hm5 subtypes: effect of G protein-coupled receptor kinases GRK2, GRK4, GRK5 and GRK6. *J Pharmacol Exp Ther* **284**:1218–1226.
- Van Koppen CJ and Jakobs KH (2004) Arrestin-independent internalization of G protein-coupled receptors. *Mol Pharmacol* **66**:365–367.
- Vickery RG and von Zastrow M (1999) Distinct dynamin-dependent and -independent mechanisms target structurally homologous dopamine receptors to different endocytic membranes. *J Cell Biol* **144**:31–43.
- Virlon B, Firsov D, Cheval L, Reiter E, Troispoux C, Guillou F, and Elalouf JM (1998) Rat G protein-coupled receptor kinase GRK4: identification, functional expression and differential tissue distribution of two splice variants. *Endocrinology* **139**:2784–2795.
- Watanabe H, Xu J, Bengra C, Jose PA, and Felder RA (2002) Desensitization of human renal D<sub>1</sub> dopamine receptors by G protein-coupled receptor kinase 4. *Kidney Int* **62**:790–798.
- Willems JM, Challiss RA, and Nahorski SR (2003) Non-visual GRKs: are we seeing the whole picture? *Trends Pharmacol Sci* **24**:626–633.
- Zeng C, Sanada H, Watanabe H, Eisner GM, Felder RA, and Jose PA (2004) Functional genomics of the dopaminergic system in hypertension. *Physiol Genomics* **19**:233–246.

---

**Address correspondence to:** Kim A. Neve, VA Medical Center (R&D-30), 3710 SW US Veterans Hospital Rd, Portland, OR 97239-2999. E-mail: nevek@ohsu.edu

---